

A NON-COMMUTATIVE ANALOGUE OF GAUSSIAN HILBERT SPACES

MIHAI POPA

ABSTRACT. The paper gives analogues of some starting results in the theory of Gaussian Hilbert Spaces for semicircular distributed random variables. The transition from the commutative to the free frame is done considering matrices of increasing dimension and utilizing the Amitsur-Levitzki Theorem.

1. INTRODUCTION

The Gaussian Hilbert spaces (i.e. complete linear spaces of random variables with centered Gaussian distributions) are referred in many works (see [5]) as an important structure in probability theory, stochastic processes and fields, stochastic integration, quantum field theory etc. Two well-know basic results in this topic are that a vector is Gaussian if and only if any linear combination of its components is also Gaussian and that any Gaussian Hilbert space can be seen as the space of sums of creation and annihilation operators on the Symmetric Fock Space over some unique, up to an isometry, Hilbert space.

This paper gives analogues of the results above for semicircular distributed random variables which play in the Voiculescu's Free Probability theory a similar role of Gaussian variables in the classical probability frame. The main idea in the proof of the classical results is identifying the scalar coefficients of some polynomials in several commuting variables. In the non-commutative frame matrices of increasing dimension are replacing the scalars and while the technique deployed many times in Free Probability theory for obtaining analogous results to classical statements uses the property of matrices with Gaussian distributed random entries and increasing dimension to asymptotically behave as free semicircular variables, here the transition from commutative to free frame is done via the Amitsur-Levitzki Theorem.

Sections 2. and 3. contain preliminary definitions and constructions, primarily recalling the ones from [11] and [8]; the main results are given in Section 3.

2. B-PROBABILITY SPACES AND B-SEMICIRCULAR RANDOM VARIABLES

Definition 2.1. Let B be a C^* -algebra. By a B -probability space, or a non-commutative probability space over B , we will understand a pair $(\mathcal{A}, \varphi)$ consisting of an algebra $\mathcal{A}$ over B and a conditional expectation $\varphi : \mathcal{A} \longrightarrow B$. If $\mathcal{A}$ is a $*$ -algebra, the pair $(\mathcal{A}, \varphi)$ will be called $*$ - B -probability space.

Consider the moment function $\hat{\mu} = (\mu^{(n)})_{n \in \mathbb{N}}$,

$$\mu^{(n)} : \underbrace{\mathcal{A} \otimes_B \mathcal{A} \otimes_B \cdots \otimes_B \mathcal{A}}_{n \text{ times}} \longrightarrow B$$

1

given by $\mu^{(n)}(a_1 \otimes \cdots \otimes a_n) = \varphi(a_1 \cdots a_n)$ and the correspondent cumulant function $\hat{k} = (k^{(n)})_{n \in \mathbb{N}}$

$$k^{(n)} : \underbrace{\mathcal{A} \otimes_B \mathcal{A} \otimes_B \cdots \otimes_B \mathcal{A}}_{n \text{ times}} \longrightarrow B$$

given by

$$(1) \quad \mu^{(n)}(a_1 \otimes \cdots \otimes a_n) = \sum_{\pi \in NC(n)} k_\pi[a_1 \otimes \cdots \otimes a_n]$$

where $NC(n)$ is the set of all non-crossing partitions on $\{1, 2, \dots, n\}$ and $k_\pi[a_1 \otimes \cdots \otimes a_n]$ is defined recursively by:

- (a) $k_{\pi_1 \cup \pi_2} = k_{\pi_1} k_{\pi_2}$, for any $\pi_1 \in NC(1, 2, \dots, n-j), \pi_2 \in NC(n-j+1, \dots, n)$
- (b) $k_{\mathbb{1}}[a_1 \otimes \cdots \otimes a_n] = k^{(n)}(a_1 \otimes \cdots \otimes a_n)$, for $\mathbb{1}$ the partition with a single equivalence class
- (c) $k_\pi[a_1 \otimes \cdots \otimes a_p \otimes a_{p+1} \otimes \cdots \otimes a_{p+r} \otimes \cdots \otimes a_n] = k^{(n_r)}(a_1 \otimes \cdots \otimes a_p \otimes k_\sigma[a_{p+1} \otimes \cdots \otimes a_{p+r}] a_{p+r+1} \otimes \cdots \otimes a_n)$, for any $\sigma \in NC(p+1, \dots, p+r)$ and π the partition obtaining by $\sigma \cup (a_1, \dots, a_p, a_{p+r}, \dots, a_n)$

For illustrations, the explicit formulas for connections between the small order moment and cumulant functions are

$$\mu^{(1)}(a_1) = k^{(1)}(a_1) = \varphi(a_1)$$

$$\begin{aligned} \mu^{(2)}(a_1 \otimes a_2) &= k_{\{(1,2)\}}[a_1 \otimes a_2] + k_{\{(1),(2)\}}[a_1 \otimes a_2] \\ &= k^{(2)}(a_1 \otimes a_2) + k^{(1)}(a_1)k^{(1)}(a_2) \end{aligned}$$

$$\begin{aligned} k^{(2)}(a_1 \otimes a_2) &= \mu^{(2)}(a_1 \otimes a_2) - \mu^{(1)}(a_1)\mu^{(1)}(a_2) \\ &= \varphi(a_1 a_2) - \varphi(a_1)\varphi(a_2) \end{aligned}$$

$$\begin{aligned} \mu^{(3)}(a_1 \otimes a_2 \otimes a_3) &= k_{\{(1,2,3)\}}[a_1 \otimes a_2 \otimes a_3] + k_{\{(1),(2,3)\}}[a_1 \otimes a_2 \otimes a_3] \\ &\quad + k_{\{(1,2),(3)\}}[a_1 \otimes a_2 \otimes a_3] + k_{\{(1,3),(2)\}}[a_1 \otimes a_2 \otimes a_3] \\ &\quad + k_{\{(1),(2),(3)\}}[a_1 \otimes a_2 \otimes a_3] \\ &= k^{(3)}(a_1 \otimes a_2 \otimes a_3) + k^{(1)}(a_1)k^{(2)}(a_2 \otimes a_3) \\ &\quad + k^{(2)}(a_1 \otimes a_2)k^{(1)}(a_3) + k^{(2)}(a_1 \otimes k^{(1)}(a_2)a_3) \\ &\quad + k^{(1)}(a_1)k^{(1)}(a_2)k^{(1)}(a_3) \end{aligned}$$

$$\begin{aligned} k^{(3)}(a_1 \otimes a_2 \otimes a_3) &= \mu^{(3)}(a_1 \otimes a_2 \otimes a_3) - \mu^{(1)}(a_1)\mu^{(2)}(a_2 \otimes a_3) \\ &\quad - \mu^{(2)}(a_1 \otimes \mu^{(1)}(a_2)a_3) - \mu^{(2)}(a_1 \otimes a_2)\mu^{(1)}(a_3) \\ &\quad + 2\mu^{(1)}(a_1)\mu^{(1)}(a_2)\mu^{(1)}(a_3) \\ &= \varphi(a_1 a_2 a_3) - \varphi(a_1)\varphi(a_2 a_3) - \varphi(a_1 \varphi(a_2)a_3) \\ &\quad - \varphi(a_1 a_2)\varphi(a_3) + 2\varphi(a_1)\varphi(a_2)\varphi(a_3) \end{aligned}$$

Given a family $(X_1, \dots, X_m)$ of elements from $\mathcal{A}$, the information about their joint moments with respect to φ is encoded in the collection of mappings

$$\xi = (\xi_{n;i_0, \dots, i_n})_{n \in \mathbb{N}; i_0, \dots, i_n \in \{1, \dots, m\}},$$

where $\xi_{n;i_0, \dots, i_n} : \underbrace{B \times \dots \times B}_n \longrightarrow B$ are given by

$$\begin{aligned} \xi_{n;i_0, \dots, i_n}(b_1, \dots, b_n) &= k^{(n+1)}(X_{i_0} \otimes b_1 X_{i_1} \otimes \dots \otimes b_n X_{i_n}) \\ &= k^{(n+1)}(X_{i_0} b_1 \otimes X_{i_1} b_2 \otimes \dots \otimes X_{i_n}) \end{aligned}$$

or, equivalently, in the collection

$$\eta = (\eta_{n;i_0, \dots, i_n})_{n \in \mathbb{N}; i_0, \dots, i_n \in \{1, \dots, m\}}$$

where

$$\begin{aligned} \eta_{n;i_0, \dots, i_n}(b_1, \dots, b_n) &= \mu^{(n+1)}(X_{i_0} \otimes b_1 X_{i_1} \otimes \dots \otimes b_n X_{i_n}) \\ &= \mu^{(n+1)}(X_{i_0} b_1 \otimes X_{i_1} b_2 \otimes \dots \otimes X_{i_n}) \end{aligned}$$

since moments are polynomials in cumulants and viceversa.

If $X_1 = X_2 = \dots = X_m = X$, we will write $\xi_{n;X}$, respectively $\eta_{n;X}$ for the functionals above.

Remark 2.1. In the case $B = \mathbb{C}$, one only needs the coefficients $\xi_{n;i_0, \dots, i_n}(1, \dots, 1)$ (respectively the coefficients $\eta_{n;i_0, \dots, i_n}(1, \dots, 1)$), since

$$\xi_{n;i_0, \dots, i_n}(b_1, \dots, b_n) = b_1 \dots b_n \xi_{n;i_0, \dots, i_n}(1, \dots, 1)$$

(and the analogous relation for η 's).

Definition 2.2. For $(\mathcal{A}, \varphi)$ a $*$ - B -probability space, a selfadjoint element x from $\mathcal{A}$ is called B -semicircular if $\xi_{n;x} = 0$ for all $n \neq 1$.

Remark 2.2. The relation (1) enable us to formulate the above definition in terms of moments: $x = x^* \in \mathcal{A}$ is B -semicircular if

$$\eta_{n;x}(b_1, \dots, b_n) = \mu^{(n+1)}(x \otimes b_1 x \otimes \dots \otimes b_n x) = \sum_{\pi \in NCPP(n)} k_\pi[x \otimes b_1 x \otimes \dots \otimes b_n x]$$

where $NCPP(n)$ denotes all the pair-partitions from $NC(n)$.

Definition 2.3. (see [1]) The semicircular elements $x_1, \dots, x_m$ from $(\mathcal{A}, \varphi)$, a non-commutative probability space, are said to satisfy the 0-Wick Theorem if all their coefficients $\xi_{p;i_0, \dots, i_n}(1, \dots, 1)$ are zero unless $p = 1$, or, equivalently, if their joint moments are computed by the formula:

$$\varphi(x_{i_1} \dots x_{i_p}) = \sum_{\gamma \in NCPP(p)} \left(\prod_{(k,l) \in \gamma} \varphi(x_{i_k} x_{i_l}) \right), \text{ for any } i_1, \dots, i_p \in \{1, \dots, m\}$$

3. ON A PARTICULAR HILBERT BIMODULE

Let $\mathcal{H}^0$ be a real Hilbert space and $\mathcal{H}$ its complexification. For $f \in \mathcal{H}^0$ we denote by $a_1(f)$, $a_1^*(f)$, $G_1(f)$, $a_0(f)$, $a_0^*(f)$, $G_0(f)$ the CCR operators of annihilation, creation and Gaussian (sum of the creation and annihilation operators) over f , respectively their full Fock Space analogues. Let Γ_j be the von Neumann Algebra generated by $\{G_j(f), f \in \mathcal{H}\}$, $j = 0, 1$. In both situations, the vacuum state will be denoted by $\langle \Omega, \cdot \Omega \rangle$.

Denote by $\mathcal{K}(l^2(\mathbb{N}))$ the C^* -algebra of compact operators on $l^2(\mathbb{N})$; for any $n \in \mathbb{N}$,

$M_n(\mathbb{C})$ will be seen as a subalgebra of $\mathcal{K}(l^2(\mathbb{N}))$ via the identification to the set of elements with non-zero entries only in the upper left $n \times n$ -corner

For $(\mathcal{A}, \varphi)$ a non-commutative probability space and $\tilde{\mathcal{A}} = \mathcal{K}(l^2(\mathbb{N})) \otimes \mathcal{A}$, (respectively $M_n(\mathcal{A}) = M_n(\mathbb{C}) \otimes \mathcal{A}$), $\tilde{\varphi} = id_{\mathcal{K}(l^2(\mathbb{N}))} \otimes \varphi$ (respectively $\varphi_n = id_{M_n(\mathbb{C})} \otimes \varphi$) the $\mathcal{K}(l^2(\mathbb{N}))$, respectively $M_n(\mathbb{C})$ -probability spaces $(\tilde{\mathcal{A}}, \tilde{\varphi})$ and $(M_n(\mathcal{A}), \varphi_n)$ will be considered with the bimodule structure given by the linear extension of the action

$$Ax = xA = A \otimes x, \text{ for any } x \in \mathcal{A} \text{ and } A \text{ in } \mathcal{K}(l^2(\mathbb{N})) \text{ or in } M_n(\mathbb{C})$$

Definition 3.1. For $\mathcal{B}$ a C^* -algebra and $\mathcal{E}$ a Hilbert $\mathcal{B}$ -bimodule, the Full Fock Bimodule over $\mathcal{E}$ is the semi-inner-product $\mathcal{B}$ -bimodule

$$\mathcal{F}(\mathcal{E}) = \mathcal{B}\tilde{\Omega} \oplus \mathcal{E} \oplus (\mathcal{E} \otimes_{\mathcal{B}} \mathcal{E}) \oplus (\mathcal{E} \otimes_{\mathcal{B}} \mathcal{E} \otimes_{\mathcal{B}} \mathcal{E}) \oplus \dots$$

equipped with the pairing

$$\langle \cdot, \cdot \rangle : \mathcal{F}(\mathcal{E}) \times \mathcal{F}(\mathcal{E}) \rightarrow \mathcal{B}$$

given by the linear extension of $(A, B \in \mathcal{B}, \tilde{x}_j, \tilde{y}_j \in \mathcal{E})$:

$$\begin{aligned} \langle A, B \rangle &= A^* B \\ \langle \widetilde{x_1} \otimes \dots \otimes \widetilde{x_k}, \widetilde{y_1} \otimes \dots \otimes \widetilde{y_l} \rangle &= \delta_k^l \langle \widetilde{x_k} \langle \widetilde{x_1} \otimes \dots \otimes \widetilde{x_{k-1}}, \widetilde{y_1} \otimes \dots \otimes \widetilde{y_{k-1}} \rangle \widetilde{y_k} \rangle \\ &= \delta_k^l \langle \widetilde{x_k}, \langle \widetilde{x_{k1}}, \dots, \langle \widetilde{x_1}, \widetilde{y_1} \rangle \dots \widetilde{y_{k-1}} \rangle \widetilde{y_k} \rangle \end{aligned}$$

The Symmetric Fock Bimodule $\mathcal{S}(\mathcal{E})$ over $\mathcal{E}$ is the submodule of $\mathcal{F}(\mathcal{E})$ generated by the elements of the form $\underbrace{\widetilde{a} \otimes \widetilde{a} \otimes \dots \otimes \widetilde{a}}_{n \text{ times}}$, $a \in \mathcal{E}$, $n \in \mathbb{N}$.

For $\tilde{x} \in \mathcal{E}$ we consider, as in [8], $G(\tilde{x}) = a(\tilde{x}) + a^*(\tilde{x})$, where $a(\tilde{x})$ and $a^*(\tilde{x})$ are respectively the annihilation and the creation operators on $\mathcal{F}(\mathcal{E})$ given by

$$\begin{aligned} a(\tilde{x})A &= 0, \quad A \in \mathcal{B} \\ a(\tilde{x})\widetilde{x_1} \otimes \dots \otimes \widetilde{x_m} &= \langle \tilde{x}, \widetilde{x_1} \rangle \widetilde{x_2} \otimes \dots \otimes \widetilde{x_m}, \quad \tilde{x}_j \in \mathcal{E} \\ a^*(\tilde{x})A &= \tilde{x}A, \quad A \in \mathcal{B} \\ a^*(\tilde{x})\widetilde{x_1} \otimes \dots \otimes \widetilde{x_m} &= \tilde{x} \otimes \widetilde{x_1} \otimes \dots \otimes \widetilde{x_m}, \quad \tilde{x}_j \in \mathcal{E} \end{aligned}$$

Denote by $\tilde{H}$ the Hilbert $\mathcal{K}(l^2(\mathbb{N}))$ -bimodule $\mathcal{K}(l^2(\mathbb{N})) \otimes H$, where $Ax = xA = A \otimes x$, for any $x \in H$ and $A \in \mathcal{K}(l^2(\mathbb{N}))$, and denote by $\tilde{H}^0$ be the set of all selfadjoint elements from $\tilde{H}$ with respect to the involution given by $A \otimes x = Ax \mapsto A^*x = A^* \otimes \bar{x}$. Let $\mathfrak{B}$ be the algebra generated by $\mathcal{K}(l^2(\mathbb{N}))$ and $\{G(\tilde{x}), \tilde{x} \in \tilde{H}^0\}$. Then $\mathfrak{B}$ with the $\mathcal{K}(l^2(\mathbb{N}))$ -valued functional $\langle \tilde{\Omega}, \cdot \tilde{\Omega} \rangle$ is an $\mathcal{K}(l^2(\mathbb{N}))$ -*-probability space and we have that:

Proposition 3.1. For any $\tilde{x} \in \tilde{H}^0$, the operator $G(\tilde{x})$ is $\mathcal{K}(l^2(\mathbb{N}))$ -semicircular.

Proof. It is enough to prove the assertion for elements of the form $\tilde{x} = \sum_{j=1}^m A_j x_j$ where $x_j \in \mathcal{H}^0$, $n \in \mathbb{N}$ and $A_j = A_j^* \in M_n(\mathbb{C})$. Then $G(\tilde{x}) = \sum_{j=1}^m A_j G_0(x_j)$, where $A_j G_0(x_j)$ denotes the operator $A_j \otimes G_0(x_j)$ acting on $\mathcal{K}(l^2(\mathbb{N})) \otimes \mathcal{F}(H)$ and for any $B_j \in \mathcal{K}(l^2(\mathbb{N}))$ one has that

$$\begin{aligned}
k^{(p+1)}(G(\tilde{x}) \otimes B_1 G(\tilde{x}) \cdots \otimes B_p G(\tilde{x})) &= \\
&= k^{(p+1)}\left(\sum_{j=1}^m A_j G(x_j) \otimes B_1 \sum_{j=1}^m A_j G(x_j) \cdots \otimes B_p \sum_{j=1}^m A_j G(x_j)\right) \\
&= \sum_{\substack{(j_0, \dots, j_p) \\ 1 \leq j_k \leq m}} A_{j_0} B_1 A_{j_1} \cdots B_p A_{j_p} k^{(p+1)}(G_0(x_{j_0}) \otimes \cdots \otimes G_0(x_{j_p})) = 0
\end{aligned}$$

unless $p = 1$, since, from [1], $G_0(x_{j_0}), \dots, G_0(x_{j_p})$ form a semicircular family. $\square$

Remark 3.1. For $(H_j, \xi_j)_{j \in \mathbb{N}}$ a family of real Hilbert spaces with distinguished unit vectors ξ_j , one has:

$$\mathcal{F}(\widetilde{H_j}) \cong \mathcal{S}(\widetilde{H_j})$$

and

$$*_j \in \mathbb{N} \mathcal{F}(\widetilde{H_j}) \cong \bigotimes_{j \in \mathbb{N}} \mathcal{F}(\widetilde{H_j})$$

where $*$ denotes the free product of Hilbert bimodules, (as considered in [12] for Hilbert spaces) with amalgamation over $\mathcal{K}(l^2(\mathbb{N}))$

Proof. Since $\mathcal{F}(\bigoplus_{j \in \mathbb{N}} \widetilde{H_j}) = *_j \in \mathbb{N} \mathcal{F}(\widetilde{H_j})$ and $\mathcal{S}(\bigoplus_{j \in \mathbb{N}} \widetilde{H_j}) = \bigotimes_{j \in \mathbb{N}} \mathcal{S}(\widetilde{H_j})$ (same arguments as in [12] and [3]), it suffices to prove that $\mathcal{F}(\widetilde{H}) = \mathcal{S}(\widetilde{H})$ for H a real Hilbert Space, i.e. $\mathcal{S}(\widetilde{H})$ contains elements of the form $Aa_1 \otimes a_2 \otimes \cdots \otimes a_n$ for arbitrary $A \in M_m(\mathbb{C}) \subset \mathcal{K}(l^2(\mathbb{N}))$ and $a_1, \dots, a_n \in H$.

Consider $\tilde{a} = AA_1a_1 + A_2a_2 + \cdots + A_na_n$, where $A_j \in M_n(M_m(\mathbb{C})) \subset \mathcal{K}(l^2(\mathbb{N}))$ are given by

$$A_j = (\alpha_{k,l}^{(j)})_{k,j=1}^n \text{ for } \alpha_{k,l}^{(j)} = \begin{cases} \delta_j^k \delta_{j+1}^l I_m & \text{if } j = 1, \dots, n-1 \\ \delta_n^k \delta_1^l I_m & \text{if } j = n \end{cases}$$

Then, for $\sigma \in S_n$ and $B_j = \begin{cases} AA_1, & \text{if } j = 1 \\ A_j, & \text{if } j = 2, \dots, n \end{cases}$ one has that

$$B_{\sigma(1)} B_{\sigma(2)} \cdots B_{\sigma(n)} = \begin{cases} A & \text{if } \sigma = 1_n \\ 0 & \text{if } \sigma \neq 1_n \end{cases}$$

$$\begin{aligned}
\underbrace{\tilde{a} \otimes \tilde{a} \cdots \otimes \tilde{a}}_{n \text{ times}} &= \sum_{\sigma \in S_n} B_{\sigma(1)} a_{\sigma(1)} \otimes B_{\sigma(2)} a_{\sigma(2)} \otimes \cdots \otimes B_{\sigma(n)} a_{\sigma(n)} \\
&= \sum_{\sigma \in S_n} B_{\sigma(1)} B_{\sigma(2)} \cdots B_{\sigma(n)} a_{\sigma(1)} \otimes \cdots \otimes a_{\sigma(n)} \\
&= Aa_1 \otimes a_2 \otimes \cdots \otimes a_n
\end{aligned}$$

$\square$

4. SEMICIRCULAR AND COMPLEX SEMICIRCULAR FAMILIES

By a Gaussian Hilbert space it will be understood a closed linear subspace in $L^2_{\mathbb{R}}(\Omega, \Sigma, P)$ where any element is a centered Gaussian random variable, for (Ω, Σ, P) a (classical) probability space. Is said that the random variables $X_1, \dots, X_n$ are jointly Gaussian or form a Gaussian family if the vector $(X_1, \dots, X_n)$ is Gaussian-distributed, or, equivalently, the joint moments of $X_1, \dots, X_n$ are computed according to the Wick Theorem (another equivalent condition is that the joint classical cumulants of $X_1, \dots, X_n$ are all zero, except the ones of order 2 (See [5])).

Consider a non-commutative probability space $(\mathcal{A}, \varphi)$; the (selfadjoint) elements $x_1, \dots, x_n \in \mathcal{A}$ are said to form a semicircular family if the vector $(x_1, \dots, x_n)$ is the limit of a (non-commutative) Free Central Limit Theorem, or equivalently, the joint moments of $x_1, \dots, x_n$ are computed according to the 0-Wick Theorem (see [11]).

The following result is the analogue of the classical theorem (cf. [5]):

The vector-valued random variable $(X_1, \dots, X_n)$ is Gaussian if and only if for any $\alpha_j \in \mathbb{R}$ ($j = 1, \dots, n$), the random variable $\sum_{j=1}^n \alpha_j X_j$ is Gaussian.

Theorem 4.1. *Let $(\mathcal{A}, \varphi)$ be a non-commutative $*$ -probability space and $x_1, \dots, x_n \in \mathcal{A}$. Then $(x_1, \dots, x_n)$ form a semicircular family if and only if $\tilde{x} = \sum_{j=1}^n A_j x_j$ is $\mathcal{K}(l^2(\mathbb{N}))$ -semicircular in $(\tilde{\mathcal{A}}, \tilde{\varphi})$, for any $A_j \in \mathcal{K}(l^2(\mathbb{N}))_{SA}$.*

Proof. Suppose $(x_1, \dots, x_n)$ form a semicircular family. Then, for $p \neq 1$, one has that:

$$\begin{aligned}
 \xi_{p, \tilde{x}}(B_1, \dots, B_p) &= k^{(p+1)}(\tilde{x} \otimes B_1 \tilde{x} \otimes \dots \otimes B_p \tilde{x}) \\
 &= k^{(p+1)}\left(\sum_{j=1}^n A_j x_j \otimes B_1 \sum_{j=1}^n A_j x_j \otimes \dots \otimes B_p \sum_{j=1}^n A_j x_j\right) \\
 &= k^{(p+1)}\left(\sum_{(j_0, \dots, j_p)} A_{j_0} x_{j_0} \otimes B_1 A_{j_1} x_{j_1} \otimes \dots \otimes B_p A_{j_p} x_{j_p}\right) \\
 &= k^{(p+1)}\left(\sum_{(j_0, \dots, j_p)} A_{j_0} B_1 A_{j_1} \dots B_p A_{j_p} x_{j_0} \otimes \dots \otimes x_{j_p}\right) \\
 &= \sum_{\substack{(j_0, \dots, j_p) \\ 1 \leq j_k \leq n}} A_{j_0} B_1 A_{j_1} \dots B_p A_{j_p} k^{(p+1)}(x_{j_0} \otimes \dots \otimes x_{j_p}) = 0
 \end{aligned}$$

since $k^{(p+1)}(x_{j_0} \otimes \dots \otimes x_{j_p}) = 0$ for $p \neq 1$, so q.e.d..

Conversely, take $\tilde{x} = \sum_{j=1}^n A_j x_j$ for $A_j \in M_m(\mathbb{C})_{SA}$. If $\tilde{x}$ is $\mathcal{K}(l^2(\mathbb{N}))$ -semicircular, then $\xi_{p, \tilde{x}}(I_m, \dots, I_m) = 0$ for $p \neq 1$, so

$$\begin{aligned}
 0 &= k^{(p+1)}(\tilde{x} \otimes \dots \otimes \tilde{x}) = k^{(p+1)}\left(\sum_{j=1}^n A_j x_j \otimes \dots \otimes \sum_{j=1}^n A_j x_j\right) \\
 &= \sum_{\substack{(j_0, \dots, j_p) \\ 1 \leq j_k \leq n}} A_{j_0} \dots A_{j_p} k^{(p+1)}(x_{j_0} \otimes \dots \otimes x_{j_p})
 \end{aligned}$$

The above expression can be seen as a polynomial $f(A_1, \dots, A_p)$ in the non-commutative variables $A_1, \dots, A_p$. Note that $f(A_1, \dots, A_p) = 0$ for any selfadjoint

$A_1, \dots, A_p \in M_m(\mathbb{C}) \subset \mathcal{K}(l^2(\mathbb{N}))$ and any $m \in \mathbb{N}$, particularly for $m = p + 1$. On the other hand, the Theorem of Amitsur-Levitzki and Lemma I.3.2 from [4] state that the lowest degree of such a polynomial with non-zero coefficients is $2(p + 1)$, therefore all the coefficients of f must be zero, i.e. $k^{(p+1)}(x_{j_0} \otimes \dots \otimes x_{j_p}) = 0$ for $p \neq 1$, so $x_1, \dots, x_n$ form a semicircular family. $\square$

Definition 4.1. *The element c from the non-commutative $*$ -probability space $(\mathcal{A}, \varphi)$ will be called complex semicircular if there are two jointly semicircular elements $s_1, s_2 \in \mathcal{A}$ such that $c = s_1 + is_2$. If s_1 and s_2 are free, then c is called circular.*

The elements $c_1, \dots, c_n \in \mathcal{A}$ are said to form a complex semicircular family if $\Re c_1, \Im c_1, \dots, \Re c_n, \Im c_n$ form a semicircular family.

For $(\mathcal{A}, \varphi)$ a $$ -B-probability space, $\tilde{c} \in \mathcal{A}$ is called complex B-semicircular if there are two B-semicircular elements $\tilde{s}_1, \tilde{s}_2 \in \mathcal{A}$ such that $\tilde{c} = \tilde{s}_1 + i\tilde{s}_2$ and for any $b_1, b_2 \in B$ if $b_1\tilde{s}_1 + b_2\tilde{s}_2$ is selfadjoint then it is B-semicircular.*

The complex semicircular elements are the free analogue of the classical complex Gaussian random variables. As in the classical case, 4.1 implies the following:

Corollary 4.2. *Let $(\mathcal{A}, \varphi)$ be a non-commutative $*$ -probability space and $c_1, \dots, c_n$ be a family of elements in $\mathcal{A}$. Then $(c_1, \dots, c_n)$ form a complex semicircular family if and only if $\tilde{x} = \sum_{j=1}^n A_j c_j$ is complex $\mathcal{K}(l^2(\mathbb{N}))$ -semicircular in $(\tilde{\mathcal{A}}, \tilde{\varphi})$, for any $A_j \in \mathcal{K}(l^2(\mathbb{N}))$.*

Let $L^2(\mathcal{F}(\tilde{H}))$ be the closure of the algebra of polynomials in $\{G(\tilde{x}), \tilde{x} \in \tilde{H}^0\}$ with coefficients in $\mathcal{K}(l^2(\mathbb{N}))$ under the norm $\tilde{x} \mapsto (\|\tilde{\varphi}(\tilde{x}^* \tilde{x})\|)^{\frac{1}{2}}$.

The following Theorem is the analogue of the classical result ([2], [5], [9]): *Let $(V, \langle, \rangle)$ be a real Hilbert space of Gaussian variables and $\mathcal{A}(V)$ be the algebra of polynomials in elements of V . Then there is a unique, up to an isometry, real Hilbert space H_0 together with an isometry $\psi : (V, \langle, \rangle) \rightarrow (\{G_1(\eta); \eta \in H_0\}, \langle \Omega, \cdot \Omega \rangle)$ which extends to an isometric isomorphism from $\overline{\mathcal{A}(V)}^{\langle \cdot, \cdot \rangle}$ to $L^2(\Gamma_1, \langle \Omega, \cdot \Omega \rangle)$.*

Theorem 4.3. *Let $(\mathcal{A}, \varphi)$ be a $*$ -non-commutative probability space, $V \subset \mathcal{A}$ a linear subspace of selfadjoint elements, and $\mathfrak{A}$ the algebra generated by the polynomials in elements of $\mathcal{K}(l^2(\mathbb{N})) \otimes V$. Then, with the above notations, we have that all the elements of $(\mathcal{K}(l^2(\mathbb{N})) \otimes V)_{SA}$ are $\mathcal{K}(l^2(\mathbb{N}))$ -semicircular if and only if there is an unique, up to an isometry, real Hilbert space $\mathcal{H}$ together with an isometry $\psi : (V, \varphi) \rightarrow (\{G_0(\eta); \eta \in \mathcal{H}\}, \langle \Omega, \cdot \Omega \rangle)$ which extends to an isometric isomorphism from $\overline{\mathfrak{A}}^{\|\tilde{\varphi}(\cdot, \cdot)\|}$ to $L^2(\mathcal{K}(\tilde{H}))$.*

Proof. Suppose we have $(\mathcal{H}, \psi)$ as in the hypothesis. Any $\tilde{x} \in \mathcal{K}(l^2(\mathbb{N})) \otimes V)_{SA}$ can be approximated with elements of the form $\sum_{j=1}^m A_j x_j$, with $x_j \in V$ and A_j in $M_n(\mathbb{C})_{SA}$ for some $n \in \mathbb{N}$.

Let $\eta_j \in \mathcal{H}$ such that $x_j = \psi^{(-1)}(G_0(\eta_j))$, $j = 1, \dots, m$. From [1], $\{G_0(\eta_j)\}_{j=1}^m$ satisfy the 0-Wick Theorem, hence so do $\{x_j\}_{j=1}^m$, therefore $(x_1 \dots x_m)$ is a semicircular family and Thm 1. implies $\tilde{x}$ is $\mathcal{K}(l^2(\mathbb{N}))$ -semicircular, q.e.d..

Conversely, suppose that all the elements from $\mathcal{K}(l^2(\mathbb{N})) \otimes V_{SA}$ are $\mathcal{K}(l^2(\mathbb{N}))$ -semicircular. For $x \in V$ let $\hat{x} = G_0^{(-1)}(\psi(x))$. If a, b are arbitrary elements from V then $\langle \hat{a}, \hat{b} \rangle = \langle \psi(a)\psi(b)\Omega, \Omega \rangle = \varphi(ab)$, therefore the uniqueness is proven.

For the existence, let V_0 be the closure of V under $\|x\| = \sqrt{\varphi(x^2)}$. Consider $\mathcal{H} = (V_0, \langle, \rangle)$ with $\langle x, y \rangle = \varphi(x^*y)$ and $x \mapsto \hat{x}$ the natural identification from V_0 to H_0 . Then $\psi(\cdot) = G_0(\cdot)$ is an isometry.

Since any selfadjoint $\tilde{x} \in \mathcal{K}(l^2(\mathbb{N})) \otimes V$ can be approximated with elements of the form $\sum_{j=1}^m A_j x_j = \sum_{j=1}^m A_j \psi^{-1}(G_0(\hat{x}_j))$ for some $n \in \mathbb{N}$, $x_1, \dots, x_m \in V$ and $A_1, \dots, A_m \in (M_n(\mathbb{C}) \otimes V)_{SA}$, from the 0-Wick theorem we have that :

$$\begin{aligned}
\langle \tilde{\Omega}, \psi(\tilde{x}_1) \dots \psi(\tilde{x}_m) \tilde{\Omega} \rangle &= \\
&= \langle \tilde{\Omega}, (\sum_{1 \leq j_1 \leq p_1} A_{1,j_1} \psi(x_{1,j_1})) \dots (\sum_{1 \leq j_m \leq p_m} A_{m,j_m} \psi(x_{m,j_m})) \tilde{\Omega} \rangle \\
&= \langle \tilde{\Omega}, \sum_{\substack{(j_1, \dots, j_m) \\ 1 \leq j_k \leq p_k}} (A_{1,j_1} \psi(x_{1,j_1})) \dots (A_{m,j_m} \psi(x_{m,j_m})) \tilde{\Omega} \rangle \\
&= \sum_{\substack{(j_1, \dots, j_m) \\ 1 \leq j_k \leq p_k}} A_{1,j_1} \dots A_{m,j_m} \langle \Omega, G_0(\widehat{x_{1,j_1}}) \dots G_0(\widehat{x_{m,j_m}}) \Omega \rangle \\
&= \sum_{\substack{(j_1, \dots, j_m) \\ 1 \leq j_k \leq p_k}} A_{1,j_1} \dots A_{m,j_m} (\sum_{\gamma \in NCPP(m)} (\prod_{(k,l) \in \gamma} \langle \Omega, G_0(\widehat{x_{k,j_k}}) G_0(\widehat{x_{l,j_l}}) \Omega \rangle)) \\
&= \sum_{\substack{(j_1, \dots, j_m) \\ 1 \leq j_k \leq p_k}} A_{1,j_1} \dots A_{m,j_m} (\sum_{\gamma \in NCPP(m)} (\prod_{(k,l) \in \gamma} \langle \widehat{x_{k,j_k}}, \widehat{x_{l,j_l}} \rangle)) \\
&= \sum_{\substack{(j_1, \dots, j_m) \\ 1 \leq j_k \leq p_k}} A_{1,j_1} \dots A_{m,j_m} (\sum_{\gamma \in NCPP(m)} (\prod_{(k,l) \in \gamma} \varphi(x_{k,j_k} x_{l,j_l}))) \\
&= \sum_{\substack{(j_1, \dots, j_m) \\ 1 \leq j_k \leq p_k}} A_{1,j_1} \dots A_{m,j_m} \varphi(x_{1,j_1} \dots x_{m,j_m}) \\
&= \sum_{\substack{(j_1, \dots, j_m) \\ 1 \leq j_k \leq p_k}} \varphi_n(A_{1,j_1} x_{1,j_1} \dots A_{m,j_m} x_{m,j_m}) \\
&= \varphi_n((\sum_{1 \leq j_1 \leq p_1} A_{1,j_1} x_{1,j_1}) \dots (\sum_{1 \leq j_m \leq p_m} A_{m,j_m} x_{m,j_m})) \\
&= \varphi_n(\tilde{x}_1 \dots \tilde{x}_m)
\end{aligned}$$

□

Corollary 4.4. *Let $(\mathcal{A}, \varphi)$ be a $*$ -non-commutative probability space, $V \subset \mathcal{A}$ a linear subspace of selfadjoint elements, and $\mathfrak{A}$ the algebra generated by the polynomials in elements of $\mathcal{K}(l^2(\mathbb{N})) \otimes V$. On $\mathcal{K}(l^2(\mathbb{N})) \otimes V$ we define the $\mathcal{K}(l^2(\mathbb{N}))$ -valued inner product $\langle \tilde{x}, \tilde{y} \rangle = \tilde{\varphi}(\tilde{x}^* \tilde{y})$. Then, $\mathcal{K}(l^2(\mathbb{N})) \otimes V$ is a inner-product $\mathcal{K}(l^2(\mathbb{N}))$ -bimodule of complex $\mathcal{K}(l^2(\mathbb{N}))$ -semicircular elements if and only if there is an unique, up to an isometry, real Hilbert space $\mathcal{H}$ together with an isometry $\psi : (V, \varphi) \rightarrow (\{G_0(\eta); \eta \in \mathcal{H}\}, \langle \Omega, \cdot \Omega \rangle)$ which extends to an isometric isomorphism from $\mathfrak{A}^{||\tilde{\varphi}(\cdot, \cdot)||}$ to $L^2(\mathcal{F}(\tilde{H}))$*

Proof. Suppose we have $(\mathcal{H}, \psi)$ as in the hypothesis. Then any $\tilde{x} \in \mathcal{K}(l^2(\mathbb{N})) \otimes V$ can be approximated in the C^* -norm by elements of the form:

$$\begin{aligned} \tilde{x}_\iota &= \sum_{j=1}^m A_j x_j, \quad \text{for some } n \in \mathbb{N}, A_j \in M_n(\mathbb{C}), x_j \in V \\ &= \sum_{j=1}^m \Re(A_j) x_j + i \sum_{j=1}^m \Im(A_j) x_j, \quad \Re(A_j), \Im(A_j) \in M_n(\mathbb{C})_{SA}, x_j \in V \\ &= \widetilde{x_1} + i \widetilde{x_2}, \quad \widetilde{x_1}, \widetilde{x_2} \in (M_n(\mathbb{C}) \otimes V)_{SA} \subset (\mathcal{K}(l^2(\mathbb{N})) \otimes V)_{SA} \end{aligned}$$

Consider now $B_1, B_2 \in \mathcal{K}(l^2(\mathbb{N}))$ such that $B_1 \widetilde{x_1} + B_2 \widetilde{x_2}$ is selfadjoint. Then

$$\begin{aligned} B_1 \widetilde{x_1} + B_2 \widetilde{x_2} &= \sum_{j=1}^m B_{1,j} x_j + \sum_{j=1}^m B_{2,j} x_j, \quad B_{1,j}, B_{2,j} \in \mathcal{F}(l^2(\mathbb{N}))_{SA}, \quad x_j \in V \\ &= \sum_{j=1}^m (B_{1,j} + B_{2,j}) \psi^{-1}(G_0((\eta_j))), \quad G_0^{-1}(\psi(x_j)) = \eta_j \in \mathcal{H} \\ &= \sum_{j=1}^m \psi^{-1}(G(B'_j \eta_j)), \quad B'_j \in \mathcal{K}(l^2(\mathbb{N}))_{SA}, \quad \eta_j \in \mathcal{H} \\ &= \psi^{-1}(G(\tilde{\eta})), \quad \tilde{\eta} = \sum_{j=1}^m B'_j \eta_j \in \widetilde{H_n^0} \end{aligned}$$

and from 4.3 we have q.e.d..

To prove that $\mathcal{K}(l^2(\mathbb{N})) \otimes V$ is an inner-product bimodule, fix $\{e_j\}_{j \in J}$ a orthonormal basis in $\mathcal{H}$. Any $\tilde{x} \in \mathcal{K}(l^2(\mathbb{N})) \otimes V$ can be approximated with as $\tilde{x}_\iota = \sum_{j=1}^m A_j \psi^{-1}(G(e_j))$, with $A_j \in \mathcal{F}(l^2(\mathbb{N}))$, therefore

$$\begin{aligned} \varphi_n(\tilde{x}^* \tilde{x}) &= id_{\mathcal{K}(l^2(\mathbb{N}))} \otimes \varphi\left(\left(\sum_{j=1}^m A_j \psi^{-1}(G(e_j))\right)^* \left(\sum_{j=1}^m A_j \psi^{-1}(G(e_j))\right)\right) \\ &= \sum_{i,j=1}^m A_i^* A_j \varphi(\psi^{-1}(G(e_i)) \psi^{-1}(G(e_j))) \\ &= \sum_{i,j=1}^m A_i^* A_j \langle e_i, e_j \rangle \\ &= \sum_{j=1}^m A_j^* A_j \geq 0, \quad \text{with equality for } \tilde{x} = 0 \end{aligned}$$

Conversely, if $\mathcal{K}(l^2(\mathbb{N})) \otimes V$ is an inner-product bimodule of $\mathcal{K}(l^2(\mathbb{N}))$ -complex semicircular elements, then $(\mathcal{K}(l^2(\mathbb{N})) \otimes V)_{SA}$ is a real linear space of $\mathcal{K}(l^2(\mathbb{N}))$ -semicirculars and the conclusion follows from 4.3. $\square$

REFERENCES

- [1] E. G. Effros; M. Popa, Feynman diagrams and Wick products associated with q -Fock space. *Proc. Natl. Acad. Sci. USA* **100** (2003), no. 15, 8629-8633
- [2] J. Glimm; A. Jaffe, Quantum Physics. A functional Integral Point of View, Second edition. *Springer-Verlag, New York*, 1987. xxii+535 pp. ISBN: 0-387-96476-2

- [3] A. Guichardet, Symmetric Hilbert spaces and related topics. Lecture Notes in Mathematics, Vol. 261. *Springer-Verlag, Berlin-New York*, 1972. v+197 pp.
- [4] N. Jacobson, PI-algebras. An introduction. Lecture Notes in Mathematics, Vol. 441. *Springer-Verlag, Berlin-New York*, 1975. iv+115 pp.
- [5] S. Janson, Gaussian Hilbert spaces. Cambridge Tracts in Mathematics, 129. *Cambridge University Press*, Cambridge, 1997. x+340 pp. ISBN: 0-521-56128-0
- [6] A. Kumjian, On certain Cuntz-Pimsner algebras. *Pacific J. Math.* **217** (2004), no. 2, 275-289.
- [7] A. Nica, D. Shlyakhtenko, R. Speicher, R -diagonal elements and freeness with amalgamation. *Canad. J. Math.* **53** (2001), no. 2, 355-381.
- [8] M. Pimsner A class of C^* -algebras generalizing both Cuntz-Krieger algebras and crossed products by $\mathbb{Z}$. Free probability theory (Waterloo, ON, 1995), 189-212, Fields Inst. Commun., 12, *Amer. Math. Soc.*, Providence, RI, 1997
- [9] B. Simon, The $P(\phi)_2$ Euclidean (quantum) field theory. Princeton Series in Physics. *Princeton University Press, Princeton, N.J.*, 1974. xx+392 pp.
- [10] R. Speicher Combinatorial theory of the free product with amalgamation and operator-valued free probability theory. *Mem. Amer. Math. Soc.* **132** (1998), no. 627, x+88 pp.
- [11] R. Speicher, Free probability theory and non-crossing partitions. Sémin. Lothar. Combin. 39 (1997), Art. B39c, 38 pp. (electronic)
- [12] D.V. Voiculescu, K. J. Dykema, A. Nica, Free random variables. CRM Monograph Series, 1. *American Mathematical Society, Providence, RI*, 1992. vi+70 pp. ISBN 0-8218-6999-X

UCLA, LOS ANGELES, CA 90095-1555

E-mail address: mvpopa@math.ucla.edu